02

ITB Syndrome

Iliotibial Band Syndrome

The classic outer-knee sting that behaves for a few kilometers, then throws a tantrum.

INJURY
Iliotibial Band Syndrome

WRITTEN BY
Liesel, Head Physio

SERIES
Injury Hub ***Top 20***

By Liesel · Head Physio · Kilometer Klub · 30+ years' experience · 11x Comrades finisher

02: INJURY HUB

Iliotibial Band Syndrome

ITB Syndrome

CAN I KEEP RUNNING?

Often yes if symptoms stay predictable and settle within 24 hours. Sharp lateral-knee pain that forces you to stop means the load is still too high.

TYPICAL TIMELINE

1 to 4 weeks for mild cases. Stubborn cases may need 6 to 8 weeks of careful rebuilding.

WHAT IT IS: IN LIESEL'S WORDS

ITB syndrome is pain on the outside of the knee caused by overload of the tissues around the distal iliotibial band. Think of the IT band like a bowstring running from your hip to the outside of your knee. When it is under too much tension or asked to absorb too many repetitions, it compresses the fat pad underneath at about 30 degrees of knee bend, right around foot strike. That compression creates the burning sting on the outer knee.

This is a load and capacity story. It tends to appear after a set distance and is especially common when runners abruptly increase volume, running downhill, add hills, or spend a lot of time on cambered roads.

> ***LIESEL'S NOTE***
> *In 30 years, I have seen more runners attack a foam roller than their hip weakness. The band is a symptom. The hips are usually the story.*

HOW IT USUALLY FEELS

The pattern is distinctive. It usually behaves for the first few kilometers, then announces itself sharply on the outer side of the knee.

- Sharp or burning pain on the outside of the knee, just above the joint line.
- Starts after a predictable distance, often getting earlier each run as it worsens.
- Downhill running and stairs are the worst provocations.
- May feel a snapping or popping sensation on the outer knee as you flex and extend.
- Tender to touch the outer knee after running.
- Settles with rest, only to return at the same point next run.

WHY RUNNERS GET THIS: REAL REASONS

It rarely appears from nowhere. Load outpaces tissue capacity and the IT band is usually the messenger, not the root cause.

- Sudden mileage or intensity spike, especially adding hills or faster running.
- Weak hip abductors allowing the knee to drop inward on each foot strike.
- Tight TFL or glutes pulling the band taut and increasing compression.
- Cambered road running, always loading the same leg as the downhill side.
- Low cadence or overstriding, increasing compression at foot strike.
- Returning to full mileage too quickly after time off.

THE BIGGEST MISTAKES I SEE

- Foam rolling the IT band directly and aggressively. It is not a muscle. You cannot stretch it out.
- Stopping completely, waiting to heal, then jumping straight back to previous mileage.
- Running through it because the first few kilometers feel fine.
- Fixing the band without ever addressing the hip weakness that caused the overload.

> **REALITY CHECK**
> *That first kilometers of feeling fine is a trap. ITB syndrome lets you warm into it, then stages a protest at exactly the same point every run.*

THE ROAD BACK TO RUNNING

The sequence matters. Returning to volume before hip capacity is the most common way runners end up back at the beginning.

1. **Reduce outer-knee irritability.**
   Temporarily cut downhill, very long runs, cambered roads, and hard sessions.
2. **Build lateral hip strength.**
   Start hip abductor work immediately. This is non-negotiable for a proper recovery.
3. **Test easy, flat running.**
   Short runs on predictable surfaces only. Assess the 24-hour response before progressing.
4. **Reintroduce hills gradually.**
   Only once flat running is consistently pain-free. Start short, stay far from maximum.
5. **Keep the strength work.**
   This injury comes back reliably when the gym work stops. That is not a coincidence.
6. **Adductor Strengthening**
   Test

> **QUICK TIP**
> *Increase your cadence by 5 to 10%. More steps per minute means less compression of the IT band at foot strike.*

REHAB PLAN: KEY EXERCISES

Hip abductor strength is the central target. These exercises are the foundation of every ITB recovery I have managed.

Clamshell Side-lying, feet together, open top knee. No hip rolling. **3 x 15/side · Daily**	**Side-Lying Leg Raise** Straight top leg, lift to 45 degrees. Heel leads. **3 x 12/side · Daily**
Lateral Band Walk Band at ankles, mini-squat, step sideways. Hips level. **3 x 12 steps/side · Daily**	**Single-Leg RDL** Hinge forward on one leg, control hip alignment. **3 x 8/side · 3x week**
Hip Drop Stand on step, drop and lift opposite hip. Pelvic control. **2 x 15/side · 3x week**	**Single-Leg Squat** Slow controlled descent. Knee tracks over second toe. **3 x 8/side · 3x week**

CROSS-TRAINING THAT WORKS

Stay fit without making it worse. These are generally well-tolerated:

- Pool running and swimming, zero impact, full cardiovascular load
- Elliptical if genuinely pain-free throughout.
- Cycling with low resistance if knee flexion angle does not provoke symptoms.
- **Avoid:** downhill running, cambered roads, and long track sessions in one direction until pain-free.

> **LIESEL'S PEN**
>
> *The IT band has a dramatic personality. It loves to behave for a few kilometers and then stage a protest. If this is your injury, please do not wait until every downhill feels like a trap. A calmer week and better hips usually beat bravado. I have seen runners take six weeks to genuinely fix this and never have it return. I have also seen runners spend two years fighting the same outer knee pain because they would not do the hip work. The choice is yours.*

INJURYHUB · QUICK REFERENCE

ITB Syndrome

Iliotibial Band Syndrome - everything you need at a glance

PAIN LOCATION

Sharp or burning pain on the outside of the knee. Starts after a predictable distance. Gets earlier each run as it worsens.

COMMON TRIGGERS

Sudden mileage increase. Downhill and cambered road running. Weak hip abductors. Tight TFL and glutes.

KEY FIX

Reduce the triggers temporarily. Start hip abductor work immediately. This is the actual fix, not foam rolling the band.

RECOVERY TIMELINE

1 to 4 weeks for mild cases. Stubborn cases need 6 to 8 weeks. The consistency of hip work determines which category you fall into.

REAL TALK: LIESEL

Some runners spend years fighting ITB syndrome because they treat the symptoms and ignore the cause. The band is not the problem. The hips are. Foam rolling feels productive. Hip work feels boring. Choose boring. The runners who fully fix this are the ones who keep doing the clamshells and lateral walks long after the pain is gone. That is the difference between solving it once and chasing it forever.

EVIDENCE BASE

A Review of Treatments for Iliotibial Band Syndrome in the Athletic Population (2013)
Effects of Conservative Treatment Strategies for ITBS in Runners (2024)
Iliotibial Band Syndrome Rehabilitation in Female Runners (2020)

KilometerKlub.com · ***Book a physio session at KilometerKlub.com/get-help***